



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P.083: THE INCREMENTAL EFFECT OF OBSTRUCTIVE SLEEP APNEA SYNDROME ON ARTERIAL STIFFNESS IN NEWLY DIAGNOSED ESSENTIAL HYPERTENSIVE SUBJECTS

K. Thomopoulos*, C. Tsioufis, A. Amfilochiou, E. Vezali, M. Alchanatis, D. Tousoulis, C. Stefanadis, I. Kallikazaros

To cite this article: K. Thomopoulos*, C. Tsioufis, A. Amfilochiou, E. Vezali, M. Alchanatis, D. Tousoulis, C. Stefanadis, I. Kallikazaros (2006) P.083: THE INCREMENTAL EFFECT OF OBSTRUCTIVE SLEEP APNEA SYNDROME ON ARTERIAL STIFFNESS IN NEWLY DIAGNOSED ESSENTIAL HYPERTENSIVE SUBJECTS, Artery Research 1:S1, S47–S47, DOI: [https://doi.org/10.1016/S1872-9312\(07\)70106-0](https://doi.org/10.1016/S1872-9312(07)70106-0)

To link to this article: [https://doi.org/10.1016/S1872-9312\(07\)70106-0](https://doi.org/10.1016/S1872-9312(07)70106-0)

Published online: 21 December 2019

P.083

THE INCREMENTAL EFFECT OF OBSTRUCTIVE SLEEP APNEA SYNDROME ON ARTERIAL STIFFNESS IN NEWLY DIAGNOSED ESSENTIAL HYPERTENSIVE SUBJECTS

K. Thomopoulos*, C. Tsioufis, A. Amfilochiou, E. Vezali, M. Alchanatis, D. Tousoulis, C. Stefanadis, I. Kallikazaros. *Department of Cardiology, Hippokraton Hospital, Athens, Greece*

Objective: Although obstructive sleep apnea syndrome (OSAS) is accompanied by increased atherosclerotic cardiovascular disease burden, its relation with arterial stiffness is not yet well determined. We investigated whether essential hypertensive subjects with OSAS are characterized by increased arterial stiffness.

Methods: Our study population consisted of 46 consecutive subjects with newly diagnosed untreated stage I-II essential hypertension suffering from OSAS (35 men, aged 49±8 years) and 53 hypertensive subjects without OSAS, matched for age, sex, and smoking status. All subjects underwent polysomnography, echocardiography and aortic stiffness evaluation by means of carotid-femoral pulse wave velocity (c-f PWV) measurements.

Results: Hypertensives with OSAS [apnea hypopnea index (AHI) ≥5] compared to hypertensives without OSAS (AHI < 5), demonstrated increased levels of body mass index (31.4±4 vs 29.3±4 kg/m², p=0.015), office systolic/diastolic blood pressure (BP) (151/99 vs 145/94 mmHg respectively, p<0.05, for both cases) and relative wall thickness (0.46±0.06 vs 0.42±0.07, p=0.010). Hypertensives with OSAS compared to those without OSAS had significantly increased c-f PWV by 9% (8.56±0.49 vs 7.85±0.93 m/s, p=0.001) and this difference remained significant even after adjustment for confounders (p=0.04). In the total study population, c-f PWV was correlated with age (r=0.35, p=0.015), office systolic BP (r=0.30, p=0.007), relative wall thickness (r=0.30, p=0.03), logAHI (r=0.389, p=0.0001) and minimum oxygen saturation (minSatO₂) (r=-0.418, p=0.0001).

Conclusions: OSAS has a significant incremental effect on aortic stiffening in the setting of middle-aged essential hypertensive subjects. This finding suggests that the presence of OSAS in a hypertensive patient accelerates vascular damage, increasing cardiovascular risk.

P.084

VARIATIONS OF INTIMA MEDIA THICKNESS DURING CARDIAC CYCLE

H. van Langen*. *University Medical Center, Nijmegen, Netherlands*

The measurement of intima media thickness (IMT) of the common carotid artery has become very popular over the last decade. The IMT is regarded as a marker for atherosclerosis. A large number of studies are conducted with IMT as study endpoint.

In this study, with two ultrasound methods, it will be shown that IMT varies with blood pressure. In the first method, a sequence of 68 common carotid artery B-mode echo images covering 3 heartbeats was stored and analyzed with dedicated software. In each image the vessel diameter and the IMT of the far wall was determined. In the second method, the RF signal of M-mode echography of the same vessel segment of the same individual was used for wall tracking technique during 3 heartbeats. The time course of the diameter of the artery and the IMT of the far wall were determined.

During diastole IMT was thicker compared to systole. The cyclic variations of IMT and diameter were out of phase. Eliminating the time course revealed that IMT and diameter correlates highly and negatively. The difference of IMT could be as large as 10% of the average IMT value. Results were explainable with incompressible material during stretching.

These results underline the importance to define at what moment during the cardiac cycle the IMT measurement is done. More accurately the absolute blood pressure on the moment IMT is determined, is of great influence. Studies that omit this information contain a methodological flaw.

P.085

ACUTE EFFECT OF VOLUME EXCHANGE ON CENTRAL HEMODYNAMICS IN PERITONEAL DIALYSIS

F. Verbeke*, W. Van Biesen, A. Pletinck, L.M. Van Bortel, R. Vanholder. *¹Ghent University Hospital, Ghent, Belgium, ²Ghent University, Ghent, Belgium*

Introduction: Although peritoneal dialysis (PD) is considered favorable regarding hemodynamic stability, little is known on potential acute effects of fluid exchange. We aimed to assess central hemodynamic changes during infusion and dwell time.

Methods: Patients were randomized to 3 different starting volumes (1000, 1500 and 2500 mL) of the non-glucose containing icodextrin, followed by addition or drainage of 200 mL every 10 minutes for 50 minutes. Hemodynamic parameters were studied using the SphygmoCor, before and after infusion of the starting volumes and after each volume change. Local

carotid systolic blood pressure (BP), augmentation index (AI) and central augmentation pressure (AP) and were used as central measures.

Results: We included 13 patients (9 male, 4 female) on PD aged 57 years (range: 40-82) and with a median BP of 126/77 mmHg (range: 94-200/51-94 mmHg). After infusion of the starting volume, central systolic BP and central AP increased by 4.7 mmHg (p=0.006) and 2.8 mmHg (p=0.005). AI increased by 5.7% (p=0.04) and heart rate decreased by 2.6 bpm (p=0.006). The difference in heart rate adjusted AI was not significant. Intraperitoneal pressure increased by 2.3 cm H₂O (p=0.03). No significant changes were observed on subsequent addition or drainage of volumes.

Conclusions: Infusion of PD fluids causes a significant acute increase in central systolic BP that persists for at least one hour and is associated with a slight decrease in heart rate. This may be due to an enhanced preload from intraperitoneal venous compression and/or to an increased wave reflection.

P.086

EVALUATION OF ARTERIAL DISTENSIBILITY IN HYPERTENSIVE PATIENTS TREATED WITH IRBESARTAN

A. Vicente Casanova^{1*}, J. Redón i Más², J. Aznar Vicente³. *¹Hypertension Unit. Clinic Hospital, Valencia, Spain, ²Hypertension Unit. Clinic Hospital, Valencia, Spain, ³Research Unit. Marina Alta Hospital, Denia-Alicante, Spain*

Introduction: The aim of the study is to evaluate the improvement of arterial distensibility by means of: (1) decrease of pulse wave velocity (PWV); (2) decrease of SBP and PP; and (3) decrease of Blood Pressure (BP), using an Angiotensin-II receptor antagonist, Irbesartan.

Material and Methods: 97 hypertensive patients, of which 43 (45.7%) were men, who attended their primary care centres two times. Mean age was 55 (SD ±10) in men and 62 (SD ±11) in women. All patients had their BP checked with a mercury sphygmomanometer, and their PWV measured with an automatic device, Complior, which registers in two different points (carotid and femoral arteries). Both measurements were taken at the initial appointment and at the six months follow up.

Appointment	SBP		DBP		CF		PP		PWV	
	A1	A2	A1	A2	A1	A2	A1	A2	A1	A2
Minimum	124	111	62	62	52	55	36	40	8	7.68
Maximum	200	176	110	94	104	104	106	95	21	21.51
Mean	157.66	140.3	92.3	81.84	77.79	79.02	65.36	58.46	12.29	11.46
SD	16.01	13.36	9.09	7.82	10.58	11.65	16.45	12.62	2.9	3.1

Results: there was a decrease in SBP, Diastolic Blood Pressure (DBP), PP, and PWV. A significant relation between the decrease of PWV and the decrease of the SBP was found.

Conclusions: (1) The PWV can be a valid indirect method to measure the arterial elasticity decrease in clinical practice, and (2) The use of drugs that act directly on the renin-angiotensin system (Irbesartan) can be useful in improving such elasticity.

P.087

PULSE PRESSURE, AUGMENTATION INDEX AND PULSE WAVE VELOCITY IN SEVERE CARDIOMYOPATHY - IMPLICATIONS FOR RISK STRATIFICATION

T. Weber^{1*}, J. Auer¹, M.F. O'Rourke², B. Eber¹. *¹Cardiology Department Klinikum, Wels, Austria, ²St. Vincent's Hospital and UNSW, Sydney, Australia*

Background: Pulse Pressure (PP), Augmentation Index (Alx), and Pulse Wave Velocity (PWV) are directly and positively associated with cardiovascular risk. However, in heart failure patients, an association between a lower PP and a worse outcome has been reported as well.

Methods: We measured Alx, using applanation tonometry (SphygmoCor) in 53 patients with severe cardiomyopathy (CMP) and 106 controls matched for age, gender, and diastolic blood pressure. All patients underwent coronary angiography for suspected coronary heart disease. In a subgroup (14 patients, 28 controls), we assessed PWV during catheter pullback.

Results: Mean age was 63.3 vs 63.6 years, ejection fraction (EF) was 28 vs 68% in patients vs controls, respectively. Central (but not peripheral) PP (32 vs 37 mmHg, p=0.01) and Alx (16 vs 23, p=0.001) were lower, ejection duration was shorter (260 vs 308 msec, p<0.00001) in patients, as compared to controls. When we subdivided the CMP patients with respect to Alx, those below the median had more advanced systolic dysfunction. In multiple regression analysis, EF was an independent predictor of Alx. PWV did not differ between patients and controls (7.8 m/s each group, p=0.99).

Conclusions: For the same level of DBP, cardiomyopathy patients show significantly lower central PP and Alx, but no difference in PWV, as compared