



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

06.05: PULSE WAVEFORM CHARACTERISTICS OR CENTRAL PRESSURE INDICES TO PREDICT ADVERSE CARDIOVASCULAR OUTCOMES IN CORONARY PATIENTS

T. Weber, M. Ammer, M. Rammer, M O'Rourke, B. Eber

To cite this article: T. Weber, M. Ammer, M. Rammer, M O'Rourke, B. Eber (2008) 06.05: PULSE WAVEFORM CHARACTERISTICS OR CENTRAL PRESSURE INDICES TO PREDICT ADVERSE CARDIOVASCULAR OUTCOMES IN CORONARY PATIENTS, Artery Research 2:3, 90–90, DOI: <https://doi.org/10.1016/j.artres.2008.08.301>

To link to this article: <https://doi.org/10.1016/j.artres.2008.08.301>

Published online: 21 December 2019

were assessed by Echo-Doppler. Early and late diastolic mitral annulus longitudinal velocities (E' and A') were assessed by tissue velocity imaging. AS was estimated by carotid-femoral pulse wave velocity (PWV).

Results: compared to controls, patients had higher ($p < 0.05$ - 0.001) PWV (9.4 ± 2 vs 8.0 ± 1 m/s) and lower transmitral E/A ratio (1.2 ± 0.5 vs 1.4 ± 0.4 , $p < 0.01$); no significant differences were found for LV midwall shortening (116 ± 13 and $119 \pm 13\%$), and EF (67 ± 6 and $69 \pm 5\%$). In the overall population, transmitral E/A correlated inversely ($p < 0.01$) with age ($r = -0.60$), SBP ($r = -0.37$), DBP ($r = -0.50$), LVMI ($r = -0.43$), and PWV ($r = -0.51$). In multivariate analysis, PWV remained independent predictor of E/A, together with age, SBP and DBP (adjusted R^2 0.43, $p < 0.05$). E'/A' showed a similar correlation pattern as E/A. **Conclusion:** in subjects free of CV disease and with preserved LV systolic function, AS affects LV diastolic filling and longitudinal myocardial velocities independently of age, LVM and BP.

doi:10.1016/j.artres.2008.08.298

06.03

COLLAGEN ACCUMULATION IS A SIGNIFICANT CONTRIBUTOR TO PULMONARY HYPERTENSION-INDUCED LARGE ARTERY STIFFENING

C.Y. Ooi, Z. Wang, N.C. Chesler.

University of Wisconsin, Madison, WI, USA

Hypoxic pulmonary hypertension (HPH) leads to large pulmonary artery (PA) stiffening, which affects right ventricular afterload. We hypothesized that vascular collagen accumulation is the major cause of large PA stiffening in HPH. The hypothesis was tested with transgenic mice that synthesize collagen type I resistant to collagenase degradation (Col1a1^{R/R}) and wild-type controls. Animals were exposed to hypoxia for 10 days and then allowed to recover. Main PAs were harvested for mechanical and biochemical tests. Effects of smooth muscle cells (SMCs) were examined using the vasoactive agents U46619 and Y27632. Pressure-diameter testing showed that left PAs of both Col1a1^{R/R} and wild-type mice stiffened with hypoxia ($p < 0.01$). Measurement of hydroxyproline biochemically confirmed that PA collagen content increased as well ($p < 0.01$). After recovery, wild-type PAs were less stiff ($p < 0.01$), whereas Col1a1^{R/R} PAs remained stiffened; biochemically, collagen content tended to decrease in both strains although less so in the Col1a1^{R/R}. Interestingly, after hypoxia, Col1a1^{R/R} PAs were less stiffened than those of wild-type mice. We speculate that differences in mean PA pressure during the 10 days of hypoxia between strains are responsible. Consistent with our hypothesis, Col1a1^{R/R} PAs also accumulated less collagen in response to hypoxia than wild-type mice. In the recovery group, stiffness and collagen content were comparable between strains. Quantitation for elastin showed no significant differences between groups. No significant differences between exposures and strains were observed in the mechanical effects of SMCs. The correlation between mechanical and biological properties suggests that collagen accumulation is critical to HPH-induced PA stiffening.

doi:10.1016/j.artres.2008.08.299

06.04

THE COMBINED EFFECT OF HIGH PULSE WAVE VELOCITY AND AUGMENTATION INDEX PREDICTS MORTALITY IN AN ELDERLY POPULATION: THE PROTEGER STUDY

A.D. Protogerou¹, M.E. Safar², T.G. Papaioannou³, P. Iaria², H. Safar², P. Ducimetier², J. Blacher².

¹ Hypertension Center, Third University Department of Medicine, Sotiria Hospital., Athens, Greece

² Paris-Descartes University, Faculty of Medicine; Hôtel-Dieu Hospital; AP-HP, Paris, France

³ 1st Department of Cardiology, Hippokraton Hospital, Athens, Greece

Objective: Carotid – femoral aortic stiffness (PWV) and pressure wave reflections (AI), both markers of cardiovascular mortality, are considered to increase in parallel. A disassociation between PWV and AI has been described. In a cohort (PROTEGER study) of old population (85.4 ± 0.5) has been previously observed that neither PWV nor AI, were predictors of mortality. The aim of the present analysis was to investigate the association between PWV and AI as well as their combined effect on mortality.

Methods: Pulse wave analysis was applied in 259 subjects for the assessment of PWV and AI. The population was divided and analyzed according to tertiles of PWV and AI.

Results: Eighty-seven subjects died during the follow up (2 years). No association was found between PWV and AI. The subjects with combined high PWV (3rd tertile) and high AI (2nd and 3rd tertiles) [group a] had significantly higher mortality compared to the rest of the population [group b] (27/56: 48.6% vs 61/203: 30.1%, $p < 0.05$, Kaplan Mayer log-rank $p = 0.003$). No significant differences

were found between group a and b regarding cardiovascular (CV) risk factors. Cox regression analysis showed that group a had higher mortality independently from age, gender, cardiovascular (CV) history and classical CV risk factors.

Conclusion: PWV and AI were not independent predictors of mortality. However, those subjects with combined increase of PWV and AI have higher mortality suggesting that: (i) the effect of PWV is more prominent in the presence of increased AI and (ii) that the disassociation between PWV and AI may be related to reduced mortality.

doi:10.1016/j.artres.2008.08.300

06.05

PULSE WAVEFORM CHARACTERISTICS OR CENTRAL PRESSURE INDICES TO PREDICT ADVERSE CARDIOVASCULAR OUTCOMES IN CORONARY PATIENTS

T. Weber¹, M. Ammer¹, M. Rammer¹, O'Rourke M², B. Eber¹.

¹ Klinikum Wels-Grieskirchen, Wels, Austria

² St Vincent's Clinic, Sydney, Australia

Background: Pulse waveform characteristics (Augmentation Index – Alx, Pressure Augmentation – AP, Pulse wave transit time – Tr) as well as central pressure indices (aortic pulse pressure – aoPP, pulse pressure amplification – PPamp, pulsatility – aortic PP/mean blood pressure) have been shown to predict cardiovascular events. We aimed to determine their relative predictive value.

Methods: We prospectively assessed Alx, Alx@75, AP, Tr as well as aoPP, PPamp and pulsatility, using radial applanation tonometry and a validated transfer function, in 520 male patients undergoing coronary angiography. Primary endpoint was a composite of all cause mortality, myocardial infarction, stroke, cardiac, cerebrovascular and peripheral revascularization. Statistics used were Cox proportional hazards regression models.

Results: Mean age was 63 years, 66.7% were hypertensives, 79.2% had CAD. During a follow up of 49.4 months, 170 patients reached the primary endpoint.

All pressure waveform characteristics and central pressure indices predicted the primary endpoint: When divided into tertiles according to the various indices, patients in the first tertile had an unadjusted HR for the combined endpoint of 0.477 (0.328-0.727) for Alx, 0.492 (0.339-0.744) for Alx@75, 0.468 (0.314-0.698) for AP, 0.435 (0.302-0.647) for Tr, 0.468 (0.331-0.685) for aoPP, 0.586 (0.406-0.858) for PPamp, and 0.585 (0.407-0.839) for pulsatility, when compared to patients in the third tertile. However, after adjustment for age, extent of coronary disease, and brachial blood pressures, only Alx, Alx@75, and Tr remained significant predictors of the combined endpoint.

Conclusion: In male patients undergoing coronary angiography, pulse waveform characteristics, but not central pressure indices, consistently and independently predict cardiovascular events.

doi:10.1016/j.artres.2008.08.301

06.06

EVIDENCE OF AUGMENTATION OF SYSTOLIC CORONARY BLOOD FLOW BY RETROGRADE WAVE REFLECTION TRAVELLING-BACK FROM THE PROXIMAL AORTA

J.E. Davies¹, Z.I. Whinnett¹, N. Hadjiilouzou¹, J. Aguado-Sierra¹, K. Parker², D.P. Francis¹, A.D. Hughes¹, J. Mayet¹.

¹ Imperial College Healthcare NHS Trust, International Centre for Circulatory Health, St Mary's Campus, Imperial College, London, United Kingdom

² Physiological Flow Unit, Imperial College, London, United Kingdom

Background: It has been proposed that blood flow in coronary arteries is augmented by waves travelling-back from the distal aorta. However, these waves have never been identified. We used wave intensity analysis to time and quantify reflected waves travelling from the proximal aorta into coronary arteries, and assess their contribution to changes in coronary blood flow with ageing and hypertension.

Method and Results: In 19 subjects (age 35-73 years) sensor-tipped intra-arterial wires were used to measure pressure and Doppler velocity in the coronary arteries and at proximal aortic root. Separated wave intensity analysis was used to identify and quantify incident and reflected waves. The reflected wave in the proximal aorta occurred in systole in all subjects (159 ± 12 ms after the ECG-R-wave) and could be seen later in the coronary arteries: left main stem (186.8 ± 10.7 ms), circumflex (188.8 ± 7.4 ms) and left anterior descending (194.4 ± 7.7 ms). In the coronary arteries the mean reflected wave represented $20.3 \pm 2.1\%$ of the incident wave magnitude, and elicited a $38.9 \pm 8.4\%$ instantaneous increase in systolic coronary blood flow velocity (0.2 to 0.28 m/s, $p < 0.002$). This augmentation was seen to increase with increasing age ($r = 0.51$, $p < 0.03$), probably