



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P1.44: EARLY DETECTION OF ATHEROSCLEROTIC DISEASE IN MILD HYPERTENSIVE PATIENTS: A STRONG REASON TO REEVALUATE CARDIOVASCULAR RISK

P. Forcada, C. Kotliar, O. Montana, C. Schulte, G. Millet, S. Obregon, J. Guerrieri, D. Turri

To cite this article: P. Forcada, C. Kotliar, O. Montana, C. Schulte, G. Millet, S. Obregon, J. Guerrieri, D. Turri (2008) P1.44: EARLY DETECTION OF ATHEROSCLEROTIC DISEASE IN MILD HYPERTENSIVE PATIENTS: A STRONG REASON TO REEVALUATE CARDIOVASCULAR RISK, Artery Research 2:3, 102–103, DOI: <https://doi.org/10.1016/j.artres.2008.08.351>

To link to this article: <https://doi.org/10.1016/j.artres.2008.08.351>

Published online: 21 December 2019

acute reduction of blood pressure by baroreceptor stimulation lowers apparent but not intrinsic stiffness in hypertensives.

doi:10.1016/j.artres.2008.08.347

P1.41

COULD MEASUREMENT OF ARTERIAL STIFFNESS PROVIDE BETTER APPROACH IN RISK ASSESSMENT THAN THE CONVENTIONAL RISK FACTOR-BASED STRATIFICATION?

B. Benczur¹, R. Bocskei², F. Molnar², M. Illyes².

¹ Hetenyi Geza County Hospital, Dept of Cardiology, Szolnok, Hungary

² TensioMed Arterial Stiffness Centre, Budapest, Hungary

Although traditional risk factors may account for 90% of the attributable cardiovascular risk their prediction of CVD is weak based on SCORE Chart. We need to find new established risk factors and to detect subclinical arterial disease to predict future coronary events. Stiffening of the aorta is one of the earliest surrogate marker of vascular damage and measurement of arterial stiffness has a growing interest in risk assessment. **Aim:** Authors investigated the correlation between the high risk state characterized by SCORE $>=5\%$ and elevated aortic pulse wave velocity (PWVao, increased arterial stiffness) measured by arteriograph.

Subject and Methods: 2243 adults were included to the analysis in which SCORE could be calculated. Sensitivity, specificity and predictive values of SCORE in detecting increased PWVao were calculated by SPSS software.

Results: Elevated PWVao ($>9,62$ m/s) was detected in 38% of patient population but sensitivity of SCORE high risk category ($>=5\%$) to detect elevated PWV was poor (33%) despite high specificity (88%) while false negative cases were in 26%. Sensitivity of SCORE was a little bit better in males (65%) but much poorer in females (17%). 10% of males and 36% of females are underestimated by SCORE assessment. The ROC curve of SCORE at the cut-off value of 5% has shown 33% sensitivity but 89% specificity.

Conclusions: If PWVao is a good surrogate of preclinical atherosclerosis SCORE risk assessment seems to be quiet acceptable in men but not in women because it markedly underestimates females CV risk.

doi:10.1016/j.artres.2008.08.348

P1.42

PRECLINICAL ATHEROSCLEROTIC DISEASE: IS IT A MARKER OF RISK OF CARDIOVASCULAR EVENTS?

P. Forcada¹, C. Kotliar¹, O. Montana², C. Schulte², S. Obregon¹, G. Millet¹, J. Guerrieri¹, D. Turri¹.

¹ Hospital Universitario Austral, Pilar, Buenos Aires, Argentina

² Clinica Dim, Buenos Aires, Argentina

Aim: We compared the severity of vascular disease (VD) by ultrasonography in patients (p.) with and without cardiovascular events (CVE), to detect high risk asymptomatic individuals.

Methods: We did in the same procedure 1) CIMT 2) Plaques characterization, 3) PWV and 4) FMD with a strict quality control. We set a score (VS) from 0 to 5 according to the severity of the VD. The CV Risk using Framingham score (FS) was also obtained from medical records.

Results: We performed a cross sectional, observational study on 581 p. (75 with CVE (AMI, Stroke, TIA, vascular thrombotic events) $62 + 10$ y.o., 73% males and 506 non CVE controls $52 + 14$ y.o. p.001, 64% males p NS). FS was high ($>20\%$) for 216p. (30,8%), moderate (10-20%) for 204p. (29%), and low ($<10\%$) for 282p. (40,2%).

Parameter	CVE (n= 75)	No CVE (n= 506)	P
SBP (mmHg)	139 ± 17	140 ± 17	NS
DBP (mmHg)	82 ± 11	85 ± 10	.03
HR (bpm)	69 ± 11	70 ± 10	NS
Left IMT(mm)	0.87 ± 0.20	0.74 ± 0.19	< 001
Right IMT (mm)	0.82 ± 0.17	0.71 ± 0.18	< 001
% abnormal IMT	32	27	NS
% Plaques	77	49	< 001
%Abnormal FMD	32	36	NS
PWV (mts / sec)	12 ± 6	10 ± 4	.02
%Abnormal PWV	34	37	NS
FS (mean)	24 ± 9	13 ± 11	< 001
VS (mean)	3,2 ± 1,3	2,3 ± 1,4	< 001

Conclusions: 1- The severity of the VD is higher in patients with CVE even when the cut off points of normality may need to be adjusted. 2- The presence of

a combination of vascular structural and functional disarrangements in asymptomatic subjects may suggest an increased risk of CVE. 3-A score of severity of VD

doi:10.1016/j.artres.2008.08.349

P1.43

FRAMINGHAM SCORE UNDERDIAGNOSES VASCULAR DISEASE IN PATIENTS UNDER CARDIOVASCULAR PREVENTION

P. Forcada¹, C. Kotliar¹, S. Obregon¹, O. Montana², C. Schulte², G. Millet¹, J. Guerrieri¹, D. Turri¹.

¹ Hospital Universitario Austral, Pilar, Buenos Aires, Argentina

² Clinica Dim, Buenos Aires, Argentina

Aim: Framingham score (FS) is used in clinical practice to estimate the CV risk of complications and is used as reference to evaluate new markers of CV risk. Recently ESH/ESC 07 guidelines stressed on the evaluation of subclinical vascular disease (VD). We analyzed, using an integrative ultrasound evaluation, the severity of VD according to increasing FS levels of risk.

Methods: We did in the same procedure 1) CIMT 2) Plaques characterization, 3) PWV and 4) FMD with a strict quality control. We set a score (VS) from 0 to 5 according to the severity of the VD. The FS was obtained from medical records.

Results: We performed a cross sectional, observational study on 702 p. (54 + 13y.o., 448(64%) males). FS was high ($>20\%$) for 216p. (30,8%), moderate (10-20%) for 204p. (29%), and low ($<10\%$) for 282p. (40,2%).

Parameter	FS Low	FS Moderate	FS High	p
Age	44 ± 12	57 ± 8	64 ± 9	< 001
Sex (% males)	53	68	74	< 001
SBP	134 ± 16	150 ± 15	145 ± 18	< 001
DBP	82 ± 10	85 ± 10	86 ± 10	< 001
Left IMT (mm)	0.66 ± 0.16	0.78 ± 0.17	0.87 ± 0.2	< 001
Right IMT (mm)	0.63 ± 0.13	0.75 ± 0.15	0.84 ± 0.2	< 001
% Plaques	30	55	31	< 001
%Abnormal FMD	32	36	44	.02
PWV (mts / sec)	9 ± 3	10 ± 3	13 ± 6	< 001
Vascular Score (mean)	1,9 ± 1,3	2,5 ± 1,3	3,1 ± 1,3	< 001

Conclusions: 1- The higher the FS, the more the severity of the VD increases. 2- Although, we have found 54,2% with a low FS with moderate to severe VS and 18% of pts. with severe VD classified as low to moderate clinical risk.

doi:10.1016/j.artres.2008.08.350

P1.44

EARLY DETECTION OF ATHEROSCLEROTIC DISEASE IN MILD HYPERTENSIVE PATIENTS: A STRONG REASON TO REEVALUATE CARDIOVASCULAR RISK

P. Forcada¹, C. Kotliar¹, O. Montana², C. Schulte², G. Millet¹, S. Obregon¹, J. Guerrieri¹, D. Turri¹.

¹ Hospital Univerdsitario Austral, Pilar, Buenos Aires, Argentina

² Clinica Dim, Buenos Aires, Argentina

Aim: To analyze the incidence and severity of subclinical vascular disease (VD) using ultrasonography in patients (p.) with essential hypertension (HT).

Methods: We did in the same procedure 1) CIMT 2) Plaques characterization, 3) PWV and 4) FMD with a strict quality control. We set a score (VS) from 0 to 5 according to the severity of the VD. The CV Risk using Framingham score (FS) was also obtained from medical records.

Results: We did a cross sectional, observational study on 604 p. (479 with stage I-II HT (ESH 07) $53,2 + 13$ y.o., 63% males and 125 normotensive NT controls $51,7 + 14$ y.o. p.003, 62% males p NS)

Parameter	HT (n= 479)	NT (n= 125)	p
SBP (mmHg)	143 ± 16	128 ± 15	< 001
DBP (mmHg)	86 ± 9	77 ± 8	< 001
HR (bpm)	70 ± 10	69 ± 9	NS
Left CIMT (mm)	0.75 ± 0.19	0.72 ± 0.10	NS
Right CIMT (mm)	0.72 ± 0.19	0.70 ± 0.19	NS
% abnormal CIMT	29	19	.02
% Plaques	49	46	NS
%Abnormal FMD	39	26	.006
PWV (mts / sec)	10,5 ± 4,3	10 ± 4	< 001
%Abnormal PWV	34	42	NS
FS (mean)	12 ± 8	8 ± 6	.001
VS (mean)	2,4 ± 1,4	2,1 ± 1,4	< 001

Conclusions: 1- Subclinical vascular disease is more frequent and severe in HT patients. 2- Indicators of subclinical vascular disease in this subset of p. allow a more precise diagnosis of subclinical vascular. (ESC/ESH guidelines 2007) 3- Vascular score of severity is higher in HT p. than in NT.

doi:10.1016/j.artres.2008.08.351

P1.45

GAMMA-GLUTAMYLTRANSFERASE - ANOTHER MARKER OF THE CARDIOVASCULAR RISK AND EARLY ARTERIAL DAMAGE?

M. Kovaite, L. Ryliskyte, Z. Petrulioniene, J. Badariene, V. Dzenkeviciute, A. Cypiene, A. Laucevicus.

Centre of Cardiology and Angiology Vilnius University Hospital Santariškių Klinikos, Vilnius, Lithuania

Purpose: The aim of the study was to evaluate whether gamma-glutamyl-transferase (γ -GT) is the marker of the cardiovascular (CV) risk assessed by the SCORE system and early arterial damage.

Methods: Serum levels of γ -GT, lipid profile, inflammatory markers and plasma glucose of 209 subjects (40-65 years old, 97 males) without clinically overt cardiovascular disease were checked. CV risk was evaluated by SCORE system. Parameters of arterial stiffness – aortic augmentation index (AIx/HR) and carotid-radial pulse wave velocity (PWV) - were obtained by applanation tonometry. Endothelium-dependent flow-mediated dilatation (FMD) in the brachial artery and carotid IMT were assessed using high resolution B-mode ultrasonography.

Results: We found that SCORE system is associated with the γ -GT. SCORE risk significantly increased with each quartile of γ -GT (ANOVA, $p=0.003$). γ -GT was significantly higher in high ($\geq 5\%$) vs. low ($< 5\%$) CV risk groups (43.82 ± 48.31 vs. 24.55 ± 22.66 , $p < 0.0001$). In the univariate analysis γ -GT was significantly associated with the FMD in the brachial artery ($p=0.025$), carotid-radial PWV ($p=0.042$) and the presence of the carotid plaques (CP) ($p=0.002$). The multiple regression analysis revealed that only gender (beta = -0.28 , $p < 0.001$), age (beta = 0.301 , $p < 0.001$) and triglycerides (beta = -0.147 , $p=0.046$) were associated with FMD. PWV was independently associated with gender (beta = -0.319 , $p < 0.001$), glucose (beta = 0.248 , $p=0.001$) and age (beta = 0.218 , $p=0.003$).

Conclusions: The study suggests that serum γ -GT is a surrogate marker of CV risk. Elevation of γ -GT is associated with the increased endothelial dysfunction, arterial stiffness and carotid plaques, but not in the independent manner.

doi:10.1016/j.artres.2008.08.352

P1.46

INTIMA-MEDIA THICKNESS VARIATION AS PREDICTOR OF ATHEROSCLEROSIS

I.M. Graf¹, F.H.B.M. Schreuder², J.M. Hameleers¹, W.H. Mess², R.S. Reneman¹, A.P.G. Hoeks¹.

¹ Maastricht University, Maastricht, Netherlands

² University Maastricht Medical Centre, Maastricht, Netherlands

Background: The aim of the present study was to ascertain in vascular diseased patients the morphological characteristics of the common carotid artery (CCA), i.e. intima-media thickness (IMT) and IMT ipsi- and bilateral intrasubject variation (Δ IMT) and to relate them to the carotid bulb stenosis degree score to establish the relevance of each parameter.

Methods: In 154 patients, referred to the vascular laboratory for CCA examination (mean 66 years, 48% male), the carotid bulb was visualized in Doppler-mode to rate the stenosis degree based on blood flow velocities. CCA IMT and Δ IMT were measured using multiple M-mode images (covering 3 heart beats). The association of morphological characteristics with stenosis degree was evaluated with Pearson correlation (r).

Results: Ipsi- and bilateral Δ IMT were stronger associated to stenosis degree score ($r=0.67$ and $r=0.60$, respectively, with $p < 0.001$) than IMT ($r=0.40$, $p < 0.001$). The averaged IMT increased slightly for stenosis degree < 5 and abruptly for the two highest values ($r=0.93$, $p=0.001$). Mean ipsi- and bilateral Δ IMT presented a consistent graded increase with stenosis degree over the entire range (0-7), reaching a correlation close to 1 ($r=0.98$ and $r=0.97$, respectively, $p < 0.001$).

Discussion: Although CCA is a region less prone to plaques, the morphological characteristics of CCA are positively correlated with stenosis degree score. The present study indicates that ipsi- and bilateral Δ IMT exhibit a stronger relation than IMT to the severity of carotid artery stenosis, underlining that Δ IMT carries different information than IMT. In conclusion, in the CCA, Δ IMT rather than IMT substantiates patient risk evaluation.

doi:10.1016/j.artres.2008.08.353

P1.47

CORRELATION BETWEEN INDICES OF ARTERIAL STIFFNESS AND ESTIMATES OF CARDIOVASCULAR EVENTS: PROMOTING CARDIOVASCULAR RISK STRATIFICATION IN CLINICAL PRACTICE

E. Malshi, C. Morizzo, F. Santini, M. Kozáková, C. Palombo.

Department of Internal Medicine, University of Pisa, Pisa, Italy

Background: Aortic and carotid stiffness are independent predictors of all cause, cardiovascular, and cerebrovascular morbidity and mortality in several clinical conditions as well as in the general population.

Aim of the study: was to investigate the associations between non-invasively determined aortic and carotid artery stiffness with vestimates of absolute cardiovascular (CV) risk in healthy subjects and subjects with risk factors.

Materials: 139 subjects were recruited [104 with one or more CV risk factors (cases: 46 males, mean age 49 years), 35 healthy controls (16 males, mean age 43 years)]. Common carotid artery stiffness was investigated by an ultrasound system with wall track option (Aloka SSD-5500, Tokyo), providing a single point local wave speed (WS). Aortic stiffness was estimated by the carotid-femoral pulse wave velocity (CF-PWV) (Complior, Paris). Ten-year absolute CV risk was estimated using both the Framingham risk score (FRS) chart and the Edinburgh University Risk Chart (EURC, including echocardiographic left ventricular hypertrophy).

Results: Cases had significantly ($p < 0.001$) higher CF-PWV (10.5 ± 2.2 vs 8.6 ± 1.3 m/s), WS (7.6 ± 1.4 vs 5.8 ± 0.8 m/s), FRS (10.8 ± 10.2 vs $3.1 \pm 3.1\%$), and EURC score (21.5 ± 16 vs $6 \pm 5\%$). Significant positive correlations ($p < 0.001$) were observed for CF-PWV and WS with FRS ($r=0.56$ and 0.52 , respectively) and EURC score ($r=0.64$ and 0.61 , respectively). When subdividing CF-PWV and WS in tertiles a significant intertertile difference was observed only for progressively increasing values of EURC score.

Conclusions: indices of subclinical cardiovascular involvement obtained by an integrated ultrasonographic investigation may help to promote CV risk stratification in clinical practice.

doi:10.1016/j.artres.2008.08.354

P1.48

AORTIC DISSECTIONS: WHY A PATIENT HAS TO WAIT (MOSCOW-BASED RETROSPECTIVE STUDY)

O. Polikina, V. Vikentyev.

Moscow State University of Medicine and Dentistry, Moscow, Russian Federation

Aortic dissections (AD) are considered among most serious life-threatening conditions requiring emergency medical intervention. However, delays in recognition and treatment often occurred.

Our aim was to determine mean time intervals between 1) first symptoms appearance (diagnostic delay, DD), 2) first contact with a doctor (Doctor-related delay, DRD) and final establishing of a confirmed diagnosis.

We analyzed 37 cases of patients admitted to general hospital with AD (84.6% males, mean age – 58.3 years).

Type A AD occurred in 58.9% cases, type B - in 41.1%. The most frequent risk factors were atherosclerosis and hypertension, 94.8%. We identified 1 case of Marfan syndrome and 1 case lues tertiarai. Thoracic pain was observed in all cases: 38% patients complained of back pain and 84,6 % of anterior chest pain. Abnormal pulsation and murmurs were found in 33.3% and 25.6% of patients respectively. The DD was 1-23 days (mean 12.4), DRD was 1- 16 days (mean 9.5). DD and DRD were significantly increased in patients of 70 years of age and older and in patients with combined pathology, including coronary heart disease ($p < 0.043$, $p < 0.14$ respectively).

Majority of the DD and DRD delays were due to either masked symptoms or blurred clinical picture, or inadequate use of existing diagnostic facilities. More attention should be paid to postgraduate continuous medical education both of GPs referring patients to a hospital and of hospital-based staff.

doi:10.1016/j.artres.2008.08.355

P1.49

EVALUATION OF ARTERIAL STIFFNESS AND CARDIOVASCULAR RISK STRATIFICATION IN A GENERAL POPULATION IN NORTHERN ITALY. THE VOBARNO STUDY

M. Salvetti, M.L. Muesan, A. Painsi, C. Monteduro, E. Belotti, C. Corbellini, G. Galbassini, C. Agabiti Rosei, L. Micheletti, C. Aggiusti, F. Marino, F. Bertacchini, Agabiti Rosei E.

University of Brescia, Brescia, Italy

Hypertension guidelines underline the importance of quantification of total cardiovascular(CV)risk;an extensive evaluation of target organ damage