



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P42: 24-HOUR CENTRAL BLOOD PRESSURE IS MORE STRONGLY ASSOCIATED TO TARGET ORGAN DAMAGE THAN BRACHIAL BLOOD PRESSURE: FIRST RESULTS OF THE VASOTENS REGISTRY


Stefano Omboni, Igor N. Posokhov, Gianfranco Parati, Vitaliy S. Barkan, Ernesto Cardona Muñoz, Elena A. Grigorieva, Irina E. Minyukhina, Maria Lorenza Muiesan, Giuseppe Mulè, Iana A. Orlova, Telmo Pereira

To cite this article: Stefano Omboni, Igor N. Posokhov, Gianfranco Parati, Vitaliy S. Barkan, Ernesto Cardona Muñoz, Elena A. Grigorieva, Irina E. Minyukhina, Maria Lorenza Muiesan, Giuseppe Mulè, Iana A. Orlova, Telmo Pereira (2017) P42: 24-HOUR CENTRAL BLOOD PRESSURE IS MORE STRONGLY ASSOCIATED TO TARGET ORGAN DAMAGE THAN BRACHIAL BLOOD PRESSURE: FIRST RESULTS OF THE VASOTENS REGISTRY, Artery Research 20:C, 67–68, DOI: <https://doi.org/10.1016/j.artres.2017.10.072>

To link to this article: <https://doi.org/10.1016/j.artres.2017.10.072>

Published online: 7 December 2019

Table	
Age category (years)	CAVI mean (SD)
<40	6.83 (0.76)
40–49	7.22 (0.86)
50–59	8.20(1.29)
60–69	8.87 (1.24)
70–79	9.60 (1.36)
80–89	11.11 (1.60)



Conclusions: This suggests CAVI is closely related to ageing and may be a useful indicator of vascular age. In initial comparisons, the slope of arterial ‘ageing’ may be steeper for Europeans, especially men over 60 years, than for Japanese and particularly Koreans, but detailed analysis has not yet been done due to lack of raw data.

References

- [1] Namekata et al. Establishing baseline criteria of cardio-ankle vascular index as a new indicator of arteriosclerosis: a cross-sectional study, *BMC Cardiovascular Disorders* 2011, 11:51
- [2] Choi SY et al. Age associated increase in arterial stiffness measured according to the cardio-ankle vascular index without blood pressure changes in healthy adults. *Journal of Atherosclerosis and Thrombosis*. Vol. 20 (2013) No. 12 p. 911–923

P41 MYOCARDIAL MECHANOENERGETIC EFFICIENCY INDEX (MMEI) AND ARTERIAL STIFFNESS: ASSOCIATION IN A GENERAL POPULATION IN NORTHER ITALY

Fabio Bertacchini, Massimo Salvetti, Anna Paini, Giulia Rubagotti, Deborah Stassaldi, Carlo Aggiusti, Giulia Maruelli, Chiara Arnoldi, Giovanni Saccà, Enrico Agabiti Rosei, Maria Lorenza Muiesan *University of Brescia, Italy*

A non-invasive approach for the estimation of mechanical efficiency through the calculation of the ratio between stroke work and HR-pressure product has been recently proposed by de Simone et al. This index, which expresses the amount of blood pumped in a single beat in 1 second by the heart, may be easily obtained by echocardiography. The aim of our study was to evaluate the determinants of myocardial mechanoenergetic efficiency index (MEEi), calculated as stroke volume/heart rate and indexed to LV mass (MEEi = MEE/LVM) in a large general population sample in Northern Italy. **Design and methods:** We evaluated 478 subjects participating in a general population study in Northern Italy (Studio Vobarno). All subjects underwent a physical examination with measurement of clinic blood pressure (BP). In all subjects laboratory examinations, 24 hours blood pressure measurement, echocardiography, and assessment of carotid-femoral pulse wave velocity (PWV) were performed.

Results: Subjects had a mean age of 58 ± 10 years, a BMI of 26 ± 4, 44% were males, 69% had arterial hypertension (55% treated). MEEi was lower in males and in patients with increased PWV. MEEi was inversely correlated with age, BMI, waist circumference, clinic and 24 hours BP, glucose, uric acid, triglycerides and directly correlated with HDL. MEEi was also inversely correlated with relative wall thickness (RWT) and PWV. At linear regression multivariate

(?) analysis MEEi remained independently related to male gender ($\beta = 0.16$, $p < 0.001$), BMI ($\beta = -0.13$, $p < 0.005$), RWT ($\beta = -0.56$, $p < 0.001$) and PWV ($\beta = -0.10$, $p < 0.05$).

Conclusions: In a large sample of general population in Northern Italy myocardial mechanoenergetic efficiency was inversely correlated with arterial stiffness, independently of multiple possible confounders.

P42 24-HOUR CENTRAL BLOOD PRESSURE IS MORE STRONGLY ASSOCIATED TO TARGET ORGAN DAMAGE THAN BRACHIAL BLOOD PRESSURE: FIRST RESULTS OF THE VASOTENS REGISTRY

Stefano Omboni ¹, Igor N. Posokhov ², Gianfranco Parati ^{3,4}, Vitaliy S. Barkan ⁵, Ernesto Cardona Muñoz ⁶, Elena A. Grigorieva ⁷, Irina E. Minyukhina ⁸, Maria Lorenza Muiesan ⁹, Giuseppe Mulè ¹⁰, Iana A. Orlova ¹¹, Telmo Pereira ¹²

- ¹Clinical Research Unit, Italian Institute of Telemedicine, Varese, Italy
- ²Hemodynamic Laboratory Ltd, Nizhny Novgorod, Russian Federation
- ³Department of Cardiology, Istituto Auxologico Italiano, Milan, Italy
- ⁴Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy
- ⁵Diagnostics Department, The Hospital within the Russian Railroad Network, Chita, Russian Federation
- ⁶University of Guadalajara, Dept. of Physiology, Guadalajara, Mexico
- ⁷South Ural State Medical University, Chelyabinsk, Russian Federation
- ⁸Volga District Medical Center, Nizhny Novgorod, Russian Federation
- ⁹Dipartimento di Scienze Mediche e Chirurgiche, Università di Brescia, Medicina 2, Spedali Civili, Brescia, Italy
- ¹⁰Unità Operativa di Nefrologia ed Ipertensione, Centro di Riferimento Regionale per l’Ipertensione Arteriosa, Policlinico Paolo Giaccone, Palermo, Italy
- ¹¹Lomonosov Moscow State University Clinic, State University, Moscow, Russian Federation
- ¹²Escola Superior de Tecnologia da Saúde de Coimbra, Instituto Politécnico de Coimbra, Coimbra, Portugal

Objective: In the present analysis of the VASOTENS study [1] baseline data, we checked whether organ damage of hypertension (TOD) i) is better associated with 24-hour central than peripheral BP and ii) is related to ambulatory arterial stiffness, estimated by pulse wave velocity (PWV) and augmentation index (Alx).

Methods: TOD in 334 hypertensive patients (mean age 53 ± 15, 52% males, 45% treated) was estimated by calculation of left ventricular mass index (LVMI), intima-media thickness (IMT) and creatinine clearance (CC). 24-hour indices were estimated through the Vasotens technology [2]. 24-hour brachial (bSBP) and aortic systolic BP (aSBP), standard deviation of bSBP, PWV and Alx were obtained. Bivariate and multivariate analysis (stepwise linear regression) was used.

Results: A significant relation was found for age, bSBP and aSBP vs. LVMI and IMT (see table). IMT was also significantly related to SBP variability and arterial stiffness, whereas age, SBP variability and Alx were significantly associated with CC. In the multivariate analysis, including all ●●●variables entered in the bivariate model, adjusted by sex, statistically significant ($p < 0.001$) association was observed for aSBP and age with LVMI (standardized regression coefficient 0.25 and 0.18, respectively), and for age with IMT (0.56) and CC (–0.53).

Correlation coefficients	LVMI (g/m ²)	IMT (mm)	CC (ml/min)
Age (years)	0.25***	0.56**	–0.53**
bSBP (mmHg)	0.23***	0.24**	–0.01
aSBP (mmHg)	0.28***	0.26**	–0.05
SD bSBP (mmHg)	0.01	0.24**	–0.19*
PWV (m/s)	0.09	0.17*	–0.14
Al (%)	0.07	0.22**	–0.18*

*** p < 0.001 ; ** p < 0.01; *p < 0.05.

Conclusions: In hypertensive patients age appears to be the major determinant of TOD, with central SBP, and marginally peripheral SBP, PWV and Alx, also playing a significant role. Our results suggest that estimation of 24-hour central hemodynamics and arterial stiffness in ambulatory conditions may help improve the individualized assessment of the BP-associated TOD.

References

- [1] Omboni S, Posokhov IN, Parati G, Avolio A, Rogoza AN, Kotovskaya YV, Mulè G, Muiesan ML, Orlova IA, Grigoricheva EA, Cardona Muñoz E, Zelveian PH, Pereira T, Peixoto Maldonado JM. Vascular Health Assessment of The Hypertensive Patients (VASOTENS) Registry: Study Protocol of an International, Web-Based Telemetering Registry for Ambulatory Blood Pressure and Arterial Stiffness. *JMIR Res Protoc* 2016;5:e137.
- [2] Omboni S, Posokhov IN, Kotovskaya YV, Protogerou AD, Blacher J. Twenty-Four-Hour Ambulatory Pulse Wave Analysis in Hypertension Management: Current Evidence and Perspectives. *Curr Hypertens Rep* 2016;18:72.

P45

RENIN AT DIFFERENT PHYSICAL ACTIVITY LEVELS IN A BI-ETHNIC POPULATION: THE AFRICAN-PREDICT STUDY

Johannes van Rooyen¹, Hugo Huisman², Leonè Malan², Catharina Mels², Ruan Kruger², Lebo Gafane-Matemane², Shani Botha², Alta Schutte²
¹Hypertension in Africa Research Team (HART) and South African Medical Research Council (SaMRC) Unit for Hypertension and Cardiovascular Disease, South Africa

²Hypertension in Africa Research Team (HART) and South African Medical Research Council (SaMRC), South Africa

Background and objectives: It is widely accepted that regular physical exercise reduces the BP, particularly in hypertensive individuals. It is recommended in the prevention of hypertension to assist in BP control. The BP lowering mechanisms of exercise remain largely elusive, we therefore evaluated the RAAS as a regulator of arterial BP.

Methods: The sub-study was embedded in the African Prospective study on the Early Detection and Identification of Cardiovascular disease and Hypertension (African- PREDICT) and included 111 white and 99 black participants aged 20–30 years.

Office- and central blood pressure as well as other cardiovascular variables were measured. Renin was analysed with an ELISA- and aldosterone with a RIA kit.

Results: The BP decreased significantly in the white but not in the black participants when tertiles for physical activity levels were compared. The total renin (renin + prorenin) level decreased significantly from 789.2 to 700.0 pg/ml ($p = 0.04$) in the white but not in the black participants (821.6 to 806.6 pg/ml; $p = 0.84$) with high physical activity level. In multiple regression analysis, in the white participants, MAP and aldosterone contributed significantly and independently to the low renin in the third percentile for physical activity level. This was not evident in the black participants.

Conclusion: Only in white participants high physical activity levels were associated with decreased blood pressure and the RAAS may be an important mechanism in this regard.

P46

ARTERIAL STIFFNESS IN RELATION TO BIRTH CHARACTERISTICS IN THE JAMAICAN 1986 BIRTH COHORT

Nadia Bennett¹, Trevor Ferguson², Novie Younger-Coleman², Marshall Tulloch-Reid², Renee Walters², Amanda Rousseau², Janeil Williams², Seeromanie Harding³, J. Kennedy Cruickshank³, Rainford Wilks²

¹King's College, London, UK

²Caribbean Institute for Health Research, The University of the West Indies, Mona, Kingston, Jamaica

³Diabetes and Nutritional Sciences Division, School of Medicine, Kings College, London, UK

Background: We tested the association between birthweight and arterial stiffness measured by aortic pulse wave velocity (PWV) and cardio-ankle vascular index (CAVI) in a birth cohort of 30 year old Jamaicans.

Methods: Participants were from the 1986 Jamaica Birth Cohort. Arterial stiffness was measured as PWV using Arteriograph 24h™ and CAVI with VaSera™ devices. Current anthropometry (height, weight, waist and hip

circumference), and brachial blood pressure measures were linked to birthweight and other early life markers of CVD risk (birth-length and maternal height). Linear regression models were used for analysis.

Results: Analyses included 235 participants 44% male, with mean \pm SD age 29.8 ± 0.7 years, birthweight 3.1 ± 0.0 kg, PWV 6.3 ± 0.1 m/s and CAVI 6.3 ± 0.1 . Bivariate models showed men had higher arterial stiffness ($p < 0.001$). Maternal height ($p = 0.031$), waist/hip ratio ($p = 0.019$), BMI ($p = < 0.001$) and blood pressure (systolic and diastolic) ($p < 0.001$) were associated with PWV, but only BMI ($p < 0.001$) was associated with CAVI. There was no association between birthweight and PWV or CAVI, $p = 0.38$ and $p = 0.41$ respectively. In multivariable models, associations between birthweight and PWV and CAVI did not change after controlling for gender, BMI, and SBP. Positive associations (coef \pm SE) between PWV and BMI (0.03 ± 0.01 $p = < 0.01$) and SBP (0.03 ± 0.01 $p = 0.001$) remained significant; as did the negative associations for BMI and CAVI (-0.04 ± 0.01 $p < 0.001$).

Conclusion: Men had higher arterial stiffness even when controlling for blood pressure and the associations of blood pressure and BMI with PWV were positive whereas and BMI with CAVI was negative. Neither arterial stiffness measure was associated with birthweight.

P47

TETRAHYDROBIOPTERIN AND MARKERS OF OXIDATIVE STRESS IN A YOUNG BI-ETHNIC POPULATION: THE AFRICAN-PREDICT STUDY

Hugo Huisman^{1,2}, Carina Mels³, Johannes Van Rooyen³, Ruan Kruger³, Carla Fourie³, Lebo Gafane³, Wayne Smith³, Alta Schutte^{3,2}

¹Hypertension in Africa Research Team, North West University, South Africa

²MRC Research Unit for Hypertension and Cardiovascular Disease, North-West University, South Africa

³Hypertension in Africa Research Team (HART), North-West University, Potchefstroom, South Africa

Background/Objectives: Tetrahydrobiopterin (BH4) is a cofactor for nitric oxide synthase (NOS). Oxidative stress, reported in black populations (1), may lead to the oxidation of BH4, the uncoupling of eNOS, decreased NO and increased superoxide levels (2,3). We compared BH4 and markers of oxidative stress and their association, between black and white cohorts.

Methods: In the African-PREDICT study, we included black ($n = 300$) and white ($N = 297$) participants (aged 20–30 years). We measured blood pressure, and determined serum levels of BH4 and markers of oxidative stress.

Results: Blacks had higher blood pressure ($p < 0.001$). In blacks the following serum levels were lower: BH4 ($p < 0.0001$), total antioxidant status (TAS) ($p < 0.0001$), glutathione peroxidase (GPx), while reactive oxygen species (ROS) ($p < 0.03$) was higher. In blacks BH4 related positively with GPx in single, partial (adjusted for socio-economic status, sex, age, BMI, GGT and cotinine) and multiple regression ($R^2 = 0.16$, $\beta = 0.17$, $p = 0.02$) and glutathione reductase (GR) ($R^2 = 0.16$, $\beta = 0.15$, $p = 0.05$). We found a negative correlation between BH4 and GPx ($R^2 = 0.07$, $\beta = -0.26$, $p = 0.0006$) in whites.

Conclusions: Higher oxidative stress levels in young blacks (increased ROS, lower TAS and GPx) could explain the low concentrations of BH4, the possible uncoupling of eNOS, resulting in higher blood pressure. The uncoupling of eNOS may explain the production of ROS and peroxynitrite and may be linked to the positive correlation of BH4 with GPx and GR found in blacks, that may lead to early vascular changes.

Reference

- Mels CMC et al. 2014. Free Radical Research. Tiefenbachere CP. 2001. American Journal of Physiology – Heart and Circulatory Physiology. Bedall JK et al. 2014. Antioxidants & Redox signalling.

Poster Session I – Hypertension

P54

SEX DIFFERENCES IN AMBULATORY CENTRAL BLOOD PRESSURE AND PULSE WAVE REFLECTIONS IN UNTREATED PATIENTS

Bernhard Hametner¹, Christopher Clemens Mayer¹, Katy Whitelegg¹, Thomas Weber², Peter Fitscha³, Siegfried Wassertheurer¹

¹Center for Health & Bioresources, AIT Austrian Institute of Technology, Vienna, Austria