



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantispress.com/journals/artres>

1.2: ASSOCIATIONS BETWEEN INDICATORS OF CARDIOVASCULAR DISEASE AND PULSE WAVE ANALYSIS AND VELOCITY: A COMPARISON OF DEVICES

Elizabeth Ellins, Lucy Lennon, Olia Papacosta, Goya Wannamethee, Peter Whincup, Julian Halcox

To cite this article: Elizabeth Ellins, Lucy Lennon, Olia Papacosta, Goya Wannamethee, Peter Whincup, Julian Halcox (2018) 1.2: ASSOCIATIONS BETWEEN INDICATORS OF CARDIOVASCULAR DISEASE AND PULSE WAVE ANALYSIS AND VELOCITY: A COMPARISON OF DEVICES, Artery Research 24:C, 67–67, DOI: <https://doi.org/10.1016/j.artres.2018.10.019>

To link to this article: <https://doi.org/10.1016/j.artres.2018.10.019>

Published online: 7 December 2019



ARTERY 18, Guimaraes Oral Presentation abstracts

Oral Session I – Epidemiology + Special Populations

1.1 PROMOTION OF ARTERIAL STIFFNESS BY CHILDHOOD CANCER AND ITS CHARACTERISTICS IN ADULT LONG-TERM SURVIVORS

Natalie Arnold¹, Hiltrud Merzenich², Arthur Wingerter³, Andreas Schluz⁴, Astrid Schneider², Jürgen H. Prochaska⁵, Sebastian Göbel⁶, Marie-Astrid Neu³, Nicole Henninger³, Marina Panova-Noeva⁵, Susan Eckerle³, Claudia Spix², Irene Schidtman², Karl J. Lackner⁷, Manfred E. Beutel⁸, Norbert Pfeiffer⁹, Thomas Münzel⁶, Jörg Faber³, Philipp S. Wild⁴

¹University Medical Center of the Johannes Gutenberg-University Mainz, Germany

²Institute for Medical Biostatistics, Epidemiology and Informatics, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

³Department of Pediatric Hematology/Oncology/Hemostaseology, Center for Paediatric and Adolescent Medicine, University Medical Center of the Johannes Gutenberg University Mainz, Mainz, Germany

⁴Preventive Cardiology and Preventive Medicine, Centre for Cardiology, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

⁵Center for Thrombosis and Haemostasis, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

⁶Center for Cardiology – Cardiology I, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

⁷Institute of Clinical Chemistry and Laboratory Medicine, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

⁸Clinic for Psychosomatic Medicine and Psychotherapy, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

⁹Department of Ophthalmology, University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

Background: Vascular alterations induced by antineoplastic treatment might be considered as a possible underlying mechanism of increased cardiovascular (CV) sequelae in childhood cancer survivors (CCS). Therefore, we thought to evaluate the changes in arterial stiffness (AS) among long-term CCS compared to the general population.

Methods: AS was assessed by digital photoplethysmography (Stiffness Index (SI); m/s) among 1,002 participant of “Cardiac and Vascular late Sequelae in long-term Survivors of childhood cancer” study, diagnosed with neoplasia prior to an age of 15 years (1980–1990). A population-based subsample from the Gutenberg Health Study (GHS) (n = 5,252) was used for comparison. All subjects underwent a comprehensive, standardized clinical examination in the same study center.

Results: Compared to the population subsample with similar age range, CCS had higher SI in multivariable linear regression analysis with adjustment for cardiovascular risk factor and comorbidities ($\beta = 0.66[0.51/0.80]$; $p < 0.0001$). Moreover, SI was varying according to tumor entity with highest values in bone tumors. Interestingly, CCS demonstrated stiffer vessels than individuals from the population even in absence of a history of chemo- or radiotherapy ($\beta = 0.56[0.16/0.96]$; $p = 0.0066$) or prevalent hypertension ($\beta = 0.66[0.50/0.81]$; $p < 0.0001$) in fully-adjusted models. Finally, a 5.2-fold [3.9; 7.0] higher prevalence of SI values exceeding age-specific, population-based reference limits was observed among CCS compared to individuals from the population.

Conclusions: This is the first study demonstrating increased AS among long-term CCS. The data suggest that AS promotion might differ in individuals with childhood cancer: Cancer development and antineoplastic treatment might be relevant determinants.

1.2 ASSOCIATIONS BETWEEN INDICATORS OF CARDIOVASCULAR DISEASE AND PULSE WAVE ANALYSIS AND VELOCITY: A COMPARISON OF DEVICES

Elizabeth Ellins¹, Lucy Lennon², Olia Papacosta², Goya Wannamethee², Peter Whincup³, Julian Halcox⁴

¹Swansea University Medical School, Swansea, UK

²UCL Department of Primary Care & Population Health, UCL, London, UK

³Population Health Research Institute, St George's University of London, London, UK

⁴Swansea University Medical School, Swansea, UK

Background: Both the Sphygmocor (S) and Vicorder (V) devices can be used for Pulse Wave Analysis (PWA) and Velocity (PWV). We investigated if there were differences in the associations between markers of Cardiovascular Disease (CVD) and PWA and PWV variables as assessed by the two devices.

Methods: 1,722 men (78.5 ± 4.7 yrs) from the British Regional Heart Study underwent PWA and PWV with the Sphygmocor and Vicorder devices. Carotid artery Intima-media Thickness (cIMT), NT-proBNP and Left Ventricle Hypertrophy (LVH) detected by ECG were also assessed. Multiple regression was used to investigate the associations between cIMT, NT-proBNP and LVH with each of the following measures, central Augmentation Pressure (cAP) Augmentation Index (cAIx) Blood Pressure (cBP) and carotid to femoral PWV.

Results: Men with data obtained from both devices were included (PWA n = 1,373, PWV n = 1,122). Following adjustment for CVD risk factors, Sphygmocor cAP, cAIx and cBP were all positively associated with LVH ($p < 0.05$), the relationship between Sphygmocor PWV and LVH was negative (OR 0.87 95% CI 0.78–0.97). There were no associations between LVH and Vicorder variables. cAP measured by both devices and cAIx by Vicorder were predictors of NT-proBNP levels (ScAP Beta 0.097 $p < 0.05$, VcAP Beta 0.153 $p < 0.001$ & VcAIx Beta 0.129 $p < 0.001$). Vicorder cBP was the only predictor of cIMT (Beta 0.057 $p < 0.05$).

Conclusion: The same measures from two devices were predictors of different indicators of CVD. Further exploration is needed to understand these differences, but device used needs to be taken into consideration when comparing findings of these variables with other published results.

1.3 PREDICTION OF CARDIOVASCULAR MORTALITY AND MORBIDITY IN THE MALMÖ DIET-CANCER COHORT FOR THE IDENTIFICATION OF HEALTHY VASCULAR AGEING, USING MARKERS OF VASCULAR STATUS

Benjamin Nilsson Wadström¹, Peter Nilsson², Abd Al-Hakim Fatehali¹, Gunnar Engström¹

¹Lund University, Department of Clinical Sciences, Skane University Hospital, Malmö, Sweden