



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P96: ASSOCIATION OF METABOLIC SYNDROME AND ITS COMPONENTS WITH ARTERIAL STIFFNESS IN GENERAL POPULATION OF THE EVA STUDY

Cristina Agudo-Conde, Leticia Gomez-Sanchez, Marta Gomez-Sanchez, Rosario Alonso-Domínguez, Natalia Sánchez-Aguadero, Cristina Lugones-Sánchez, Jesus Gonzalez-Sanchez, Sara Mora-Simon, Jose I. Recio-Rodriguez

To cite this article: Cristina Agudo-Conde, Leticia Gomez-Sanchez, Marta Gomez-Sanchez, Rosario Alonso-Domínguez, Natalia Sánchez-Aguadero, Cristina Lugones-Sánchez, Jesus Gonzalez-Sanchez, Sara Mora-Simon, Jose I. Recio-Rodriguez (2018) P96: ASSOCIATION OF METABOLIC SYNDROME AND ITS COMPONENTS WITH ARTERIAL STIFFNESS IN GENERAL POPULATION OF THE EVA STUDY, Artery Research 24:C, 106–107, DOI: <https://doi.org/10.1016/j.artres.2018.10.149>

To link to this article: <https://doi.org/10.1016/j.artres.2018.10.149>

Published online: 7 December 2019

¹Cardiff University, Cardiff, Wales, UK

²Cardiff Metropolitan University, Cardiff, Wales, UK

Background: Chronic obstructive pulmonary disease (COPD) is an inflammatory condition associated with increased cardiovascular risk. COPD patients have increased aortic stiffness and increased risk of stroke. In addition, increased mid cerebral artery pulsatility index (MCAPI) is associated with increased arterial stiffness and risk of stroke in patients with Diabetes. However, the mechanisms relating to the increased risk of stroke in COPD remain unclear. Therefore, this study aims to investigate the relationship between aortic pulse wave velocity (aPWV) and the MCAPI in COPD patients.

Methods: This Cross-sectional evaluation included 20 COPD patients (mean \pm SD, age 69.0 ± 6.3 years) from the ACRADe study. aPWV was measured using the SphygmoCor system and MCAPI using transcranial Doppler ultrasound. In addition, forced expiratory volume in the first second/ forced vital capacity (FEV1/FVC) was measured using spirometry and quality of life was assessed using the St George's Respiratory Questionnaire (SGRQ). Measures of frailty were assessed using Time-Up-and-Go test (TUG) and the Comprehensive Geriatric Assessment (CGA).

Results: MCAPI was significantly associated with aPWV ($r = 0.518$, $P = 0.033$). aPWV was significantly associated with SGRQ disease impact ($r = 0.604$, $P = 0.010$) and also associated with TUG ($r = 0.561$, $P = 0.019$) and CGA ($r = 0.639$, $P = 0.006$).

Conclusions: These pilot data highlight the association between increased aortic stiffness and increased pulsatile flow velocity transmitted to the cerebral circulation of COPD patients. In addition, COPD symptoms, impact and frailty are both associated with increased aortic stiffness. However, more research is needed to investigate cause and effect between COPD symptoms, impact and its relationship between aPWV and cerebral flow pulsatility.

P94

EVALUATING CENTRAL PRESSURE IN PATIENTS WITH ACUTE ISCHEMIC STROKE IN ACUTE PHASE: PROGNOSIS AND OUTCOME

Ana Costa ¹, David Paiva ¹, Filipa Gonçalves ², Ana Campos ³, Pedro Cunha ², Jorge Cotter ⁴

¹Hospital Sra. da Oliveira, Guimarães, Portugal

²Hospital da Sra. Oliveira, Guimarães, Portugal

³Hospital da Sra. Oliveira, Guimarães, Portugal

⁴Hospital da Sra. da Oliveira, Guimarães, Portugal

Acute ischemic stroke (AIS) is defined as sudden onset of a neurologic deficit. It's the cause of about 85% of all strokes and the deficits last for more than 24 h. (1) Blood pressure (BP) is elevated in 75% or more of patients with acute stroke and different levels of peripheral BP at onset are associated with poor outcomes. In patients with AIS, management of blood pressure is still a matter of debate. Brachial pressure is a poor surrogate for aortic pressure and recent evidence suggests that central pressure is more strongly related to future cardiovascular events. In this pilot study we aimed to evaluate central pressure (CBP) in patients admitted with AIS in the acute phase (first 24 h). We evaluated 34 patients with a mean age of 72,7 years. Patients presented a mean NIHSS score 5,4 at admission (0-18) and NIHSS of 4 at discharge. Pre-AIS Rankin mean was 1 and at discharge was 2,1. Brachial systolic and diastolic blood pressures varied between 108 – 250 and 42–131 mmHg accordingly, with mean values of 147,48/78,21. Central BP varied from 102,5 – 215 mmHg systolic and 44–128,5 mmHg diastolic with mean value of 136,65/80,56 mmHg. In this sample, low values of both central and peripheral BP were associated with poor outcome (Rankin scale). This is an ongoing study aiming to evaluate central hemodynamic parameters in acute phase of AIS and at long term. The main goal is to enlarge our sample so we can be able to extract more and stronger data.

Poster Session II – Epidemiology

P95

BLOOD PRESSURE VARIABILITY, ARTERIAL STIFFNESS AND ARTERIAL REMODELING – THE MAASTRICHT STUDY

Tan Lai Zhou ^{1,2}, Ronald Henry ^{3,4,5}, Coen Stehouwer ^{6,7}, Thomas van Sloten ^{8,9,10}, Koen Reesink ^{7,11}, Abraham Kroon ^{6,7}

¹Dept. of Internal Medicine, Maastricht University, Maastricht, the Netherlands

²Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, the Netherlands

³Dept. of Internal Medicine, Maastricht University Medical Centre+, Maastricht, the Netherlands

⁴Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, the Netherlands

⁵Heart and Vascular Centre, Maastricht University Medical Centre +, Maastricht, the Netherlands

⁶Department of Internal Medicine, Maastricht University Medical Centre +, Maastricht, the Netherlands

⁷Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, the Netherlands

⁸Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, Maastricht, the Netherlands

⁹Université Paris Descartes, Sorbonne Paris Cité, Faculté de Médecine, Paris, France

¹⁰INSERM, UMR-S970, Paris Cardiovascular Research Center, Department of Epidemiology and Department of Arterial Mechanics, Paris, France

¹¹Department of Biomedical Engineering, Maastricht University, Maastricht, the Netherlands

Greater very short- to mid-term blood pressure variability (BPV) has been associated with an increased CVD risk, especially stroke. However, this link remains incompletely understood. We hypothesized that increased arterial stiffness and maladaptive carotid arterial remodeling may underlie this association. We therefore investigated the association between very short- to mid-term systolic BPV, aortic and carotid stiffness and carotid arterial remodeling using cross-sectional data from The Maastricht Study (aged 60 ± 8 years; 53% men). Aortic (carotid-femoral pulse wave velocity, $n = 1671$) and carotid stiffness (ultrasonography, $n = 1690$) were assessed. A composite index of systolic BPV was derived by standardizing and averaging systolic within-visit, 24-hour, and 7-day BPV. We performed linear regression analyses with adjustment for age, sex, glucose metabolism status, mean arterial pressure and cardiovascular risk factors. A 1-SD greater systolic BPV was statistically significantly associated with 0.10 m/s (95%CI: 0.01 – 0.20) greater cfPWV, but not with carotid distensibility ($-0.033 \cdot 10^{-3}$ /kPa [$-0.255 - 0.190$]). In addition, a 1-SD greater systolic BPV was statistically significantly associated with greater carotid circumferential wall tension (0.84 dyne/cm [$0.51 - 1.17$]), circumferential wall stress (0.79 kPa [$0.031 - 1.27$]) and intima-media thickness (8.6 μ m [$1.0 - 16.3$]). These results are indicative of maladaptive carotid remodeling, as circumferential wall tension and stress were not normalized despite greater intima-media thickness. In conclusion, greater very short- to mid-term BPV is associated with greater aortic stiffness and maladaptive carotid arterial remodeling, but not with carotid stiffness. These findings may explain, at least partially, the increased BPV-associated CVD risk, in particular stroke.

P96

ASSOCIATION OF METABOLIC SYNDROME AND ITS COMPONENTS WITH ARTERIAL STIFFNESS IN GENERAL POPULATION OF THE EVA STUDY

Cristina Agudo-Conde ¹, Leticia Gomez-Sanchez ¹, Marta Gomez-Sanchez ¹, Rosario Alonso-Dominguez ², Natalia Sánchez-Aguadero ², Cristina Lugones-Sánchez ², Jesus Gonzalez-Sanchez ², Sara Mora-Simon ², Jose I. Recio-Rodriguez ²

¹Institute of Biomedical Research of Salamanca (IBSAL), Primary Health Care Research Unit, La Alamedilla Health Center, Salamanca, Spain

²Institute of Biomedical Research of Salamanca (IBSAL), Primary Health Care Research Unit, La Alamedilla Health Center, Salamanca, Spain

Objective: The aim of this study is to investigate the relationship of Metabolic Syndrome (MetS) and its components with arterial stiffness in general population without cardiovascular diseases.

Design and method: Cross-sectional study of general population without cardiovascular diseases. There were included 500 subjects between 35–75 years old, selected by random sampling with replacement, stratified by age and gender groups using the Sanitary Card base of 5 urban health centers.

Measurements: Pulse wave velocity femoral carotid (cfPWV) was determined using the SphygmoCor System, Cardio Ankle Vascular Index (CAVI) using the VaSera. MetS was defined based on the Joint Scientific Statement National Cholesterol Education Program III.

Results: Mean age was 55.9 ± 14.2 years, mean cfPWV was 6.5 ± 2.0 m/sec and mean CAVI was 8.0 ± 1.4 . MetS was found in 14% of the subjects. cfPWV and CAVI were higher in subjects with MetS. All MetS components, except reduced HDL-cholesterol, were correlated with cfPWV and CAVI. With

correlation coefficient between $r = 0.450$ (cfPWV and diastolic blood pressure) and $r = 0.128$ (between CAVI and triglycerides). After adjustment for age and sex the correlation remains the same with the cfPWV. However, it is only maintained with the CAVI only with blood pressure. Subjects with MetS have odds ratio (OR) for both cfPWV ≥ 10 m/sec (OR = 1.884, 95 % CI 0.996–3.486) and CAVI ≥ 9 (OR = 1.810, 95 % CI 0.749–4.372).

Conclusions: The cfPWV showed the positive correlation, after adjusting it for age and sex with all the components of the MetS, however the CAVI showed the positive correlation with the arterial pressure.

P97

FAMILY PATTERNS OF CENTRAL HAEMODYNAMICS ACROSS THREE GENERATIONS IN THE MALMÖ OFFSPRING STUDY

Peter Nilsson¹, Erik Petersson Rosberg²

¹Lund University, Department of Clinical Sciences, Skane University Hospital, Jan Waldenströms gata 15, Level 5, S-20502, Malmö, Sweden

²Lund University, Department of Clinical Sciences, Skane University Hospital, Malmö, Sweden

Background: Markers of central haemodynamics have in recent years emerged as promising predictors of cardiovascular disease (CVD). Central haemodynamics are affected early in the development of vascular aging and affect organs directly attached to large arteries. Carotid-Femoral pulse wave velocity (c-f PWV), Augmentation index (Aix), and central systolic blood pressure (cSBP) are variables from indirect measurements that reflect central haemodynamic and arterial stiffness. Family patterns exist [1].

Aim: To investigate if a relationship exists for patterns of central haemodynamics across three related generations, especially c-f PWV.

Methods: In all, 1131 participants from Malmö Diet Cancer Study (MDCS) and Malmö Offspring Study (MOS) were included in this study. c-f PWV was measured in grandparents and in all offspring. Correlation analyses of c-f PWV between offspring and c-f PWV in parents and grandparents were conducted. Parents and grandparents were divided in quartiles by c-f PWV and offspring c-f PWV, and cSBP means were compared with one-way ANOVA analyses. Multiple regression analyses were conducted to adjust for age, sex, BMI, SBP and fasting glucose.

Results: c-f PWV in grandchildren was positively correlated with c-f PWV in parents ($r = 0.26$, $p < 0.001$) and in grandparents ($r = 0.29$, $p < 0.001$). Offspring c-f PWV correlated significantly with parental Aix and cSBP. Parents with high c-f PWV had offspring with statistically significant higher means of c-f PWV and cSBP than parents with low c-f PWV.

Conclusion: Measures of central haemodynamic are positively correlated across three generations in a population-based study.

References

1. Fatehali AA, Gottsäter M, Nilsson PM. Family history of cardiometabolic diseases and its association with arterial stiffness in the Malmö Diet Cancer cohort. *J Hypertens.* 2017 Nov;35(11):2262-2267.

P98

AGE AND GENDER DIFFERENCES IN VARIABILITY OF WAVE REFLECTIONS OVER 24 HOURS: THE INTERNATIONAL 24-HOUR AMBULATORY AORTIC BLOOD PRESSURE CONSORTIUM (I24ABC)

Thomas Weber¹, Siegfried Wasserheuer², James Sharman³, Cristina Giannatasio⁴, Piotr Jankowski⁵, Yan Li⁶, Alessandro Maloberti⁴, Barry McDonnell⁷, Carmel McEniery⁸, Maria Lorenza Muisan⁹, Janos Nemcsik¹⁰, Anna Paini⁹, Enrique Rodilla¹¹, Ian Wilkinson¹², Robert Zweiker¹³, Athanase Protogerou¹⁴

¹Cardiology Department, Klinikum Wels-Grieskirchen, Wels, Austria

²Austrian Institute of Technology, Vienna, Austria

³Menzies Institute for Medical Research, University of Tasmania, Australia

⁴University Milano-Bicocca, Milan, Italy

⁵Jagellonian University Krakow, Krakow, Poland

⁶Shanghai Institute of Hypertension, Shanghai Jiaotong University School of Medicine, Shanghai, China

⁷Cardiff Metropolitan University, Cardiff, UK

⁸Cambridge University, Cambridge, UK

⁹Università di Brescia, Italy

¹⁰Semmelweis University Budapest, Hungary

¹¹Valencia (Hospital de Sagunto) Universidad CEU Cardenal Herrera, Spain

¹²Cambridge University, UK

¹³Cardiology Department, Medical University Graz, Austria


¹⁴Department of Pathophysiology Medical School National and Kapodistrian University of Athens, Athens, Greece

Background: Wave reflection parameters predict cardiovascular events, but 24-hour profiles in large samples of healthy adults are unknown.

Methods: In 1645 individuals free from antihypertensive drugs from 11 centers in Europe and Asia, 24-hour blood pressure monitoring with a validated oscillometric brachial cuff (Mobilograph, I.E.M., Stolberg; Germany) was performed. Brachial waveforms were acquired and processed with ARCSolver algorithms to derive information relating to wave reflections using pulse waveform analysis (heart-rate corrected augmentation index-Alx75, augmentation pressure-AP) and wave separation analysis (backward wave amplitude-Pb, reflection magnitude-RM). Nighttime/daytime difference (N/D) was nighttime (01.00–06.00) minus daytime (09.00–21.00) values/daytime values. Participants were categorized as young (13–39 years; male/female: 219/112), middle-aged (40–66 years; male/female: 545/553), and old (67–104 years; male/female: 86/130).

Results: 24-hour measures of wave reflections increased with increasing age and were significantly lower in men compared to women (Alx75: 18.3 vs 28.0 %, AP: 10.1 vs 14.9 mm Hg, Pb: 18.9 vs 20.0 mm Hg, RM: 63.0 vs 66.2). Alx75 was higher during daytime compared to nighttime (23.3 vs 21.3%), but only in young and middle-aged participants. For all participants, AP (11.6 vs 14.5 mm Hg), Pb (18.5 vs 21.7 mm Hg), and RM (62.9 vs 68.8) were higher during nighttime compared to daytime. N/D varied with age and was more pronounced in younger individuals.

Conclusion: 24-hour variability of wave reflection parameters differs according to age and gender. In future, this information could be useful for tailoring individual cardiovascular risk management.



P99

STUDY ON THE PREVALENCE AND DETERMINANTS OF EARLY VASCULAR AGEING IN A COMMUNITY PHARMACY SETTING – PRELIMINARY RESULTS: FROM THE ASINPHAR@2ACTION (ARTERIAL STIFFNESS IN THE PHARMACIES TO (2) ACTION) PROJECT

Telmo Pereira¹, Ema Paulino², Mariana Rosa², Ana Pinto², Maria Teixeira², Patrícia Soares², Sofia Maximiano², Johannes Risse³, Sonja Gose²

¹Polytechnic Institute, Coimbra Health School, Portugal

²Holon Pharmacies, Portugal

³I.E.M, GmbH, Germany

Objective: The ASINPHAR@2action programme aims at raising awareness to early vascular ageing (EVA) through a community-based intervention. This preliminary analysis is focused on the analysis of the proportion of participants with abnormal arterial stiffness (AS) and the definition of its main determinants.

Design and method: This preliminary analysis is a cross-sectional, observational, descriptive, non-interventional study of participants enrolled in 11 communal pharmacies in Portugal (HOLON pharmacies), between April and November 2017. Blood pressure (BP) and arterial function parameters were measured with a non-invasive validated device (MOBIL-O-GRAPH, IEM®). Clinical and demographic information was gathered.