



Artery Research

ISSN (Online): 1876-4401

ISSN (Print): 1872-9312

Journal Home Page: <https://www.atlantis-press.com/journals/artres>

P172: ROLE OF ADIPOSE TISSUE AND SKELETAL MUSCLE IN MACROVASCULAR ATHEROSCLEROTIC OCCLUSIVE DISEASE-PERIPHERAL ARTERIAL DISEASE AND CAROTID ARTERY DISEASE

Joana Ferreira

To cite this article: Joana Ferreira (2018) P172: ROLE OF ADIPOSE TISSUE AND SKELETAL MUSCLE IN MACROVASCULAR ATHEROSCLEROTIC OCCLUSIVE DISEASE-PERIPHERAL ARTERIAL DISEASE AND CAROTID ARTERY DISEASE, Artery Research 24:C, 132–132, DOI: <https://doi.org/10.1016/j.artres.2018.10.225>

To link to this article: <https://doi.org/10.1016/j.artres.2018.10.225>

Published online: 7 December 2019

59. Yazici H, Tuzun Y, Pazarli H, et al. Influence of age of onset and patient's sex on the prevalence and severity of manifestations of Behçet's syndrome. *Ann Rheum Dis*. 1984; 43(6): 783–789.
60. Nihal S, Kutlubay EZ, Ucar D, Hatemi I, Uygunoglu U, Siva A, Hatemi G. Behçet's syndrome: providing integrated care. *J Multidiscip Healthc*. 2017; 10: 309–319.
61. DGS (2016). A saúde dos portugueses 2016. Lisboa: Portugal. Direção-Geral da Saúde. ISSN: 2183-5888.
62. Lopes C et al. (2017). Inquérito alimentar nacional e de atividade física. IAN-AF, 2015-2016. Porto: Universidade do Porto.
63. Instituto Nacional de Estatística. Inquérito Nacional de Saúde. 2014.
64. Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults. Report of the national cholesterol education program (NCEP). Executive Summary of The Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA*. 2001; 285: 2486-2497.
65. Malik S, Wong ND, Franklin SS, et al. Impact of the metabolic syndrome on mortality from coronary heart disease, and all cause in United States adults. *Circulation*. 2004; 110: 1245-1250.
66. Cuspidi C, Sala C, Tadic M, Gherbesi E, Grassi G, Mancia G. Association of metabolic syndrome with carotid thickening and plaque in the general population: A meta-analysis. *J Clin Hypertens (Greenwich)*. 2017; 1-7.
67. Alecu C, Gueguen S, Aubry C, Salvi P, Perret-Guillaume C, Ducrocq X, Vespignani P, Benetos A. Determinants of arterial stiffness in an apparently healthy population over 60 years. *Journal of Human Hypertension*. 2006; 20: 749–756.
68. Popele NM, Grobbee DE, Bots ML, Asmar R, Topouchian J, Reneman RS, Hoeks A, Kuip D, Hofman A, Witteman J. Association Between Arterial Stiffness and Atherosclerosis. The Rotterdam Study. *AHA Journal-Stroke*. 2001; 32: 454-460.
69. Sehestedt T, Jeppesen J, Hansen TW, Wachtell K, Ibsen H, Torp-Pedersen C, Hildebrandt P, Olsen MH. Risk prediction is improved by adding markers of subclinical organ damage to SCORE. *Eur Heart J*. 2010; 31(7): 883-91.
70. Wang F, Ye P, Luo L, Xiao W, Qi L, Bian S, et al. Association of serum lipids with arterial stiffness in a population-based study in Beijing. *Eur J Clin Invest*. 2011; 41(9): 929-36.
71. Safar ME, Czernichow SB, Blacher J. Obesity, arterial stiffness, and cardiovascular risk. *J Am Soc Nephrol*. 2006; 17(4 Suppl 2): S109-11.
72. Mäki-Petäjä KM et al. Rheumatoid Arthritis is associated with increased Aortic Pulse-Wave Velocity which is reduced by Anti-Tumor Necrosis Factor Therapy. *Circulation* AHA. 2006; 114: 1185-1192.
73. Kürüm T, Yildiz M, Soy M, Ozbay G, Alimgil L, Tüzün B. Arterial distensibility as determined by carotid-femoral pulse wave velocity in patients with behçet's disease. *Clin Rheumatol*. 2005; 24: 134-8.
74. Caldas CA, Borba EF, Bortolotto LA, Medeiros DM, Bonfa E, Gonçalves CR. Increased arterial stiffness assessed by pulse wave velocity in behçet's disease and its association with the lipid profile. *J Eur Acad Dermatol Venereol*. 2013; 27: 454-9.
75. Chang HK, Kim SK, Lee SS, Rhee MY. Arterial stiffness in behçet's disease: increased regional pulse wave velocity values. *Ann Rheum Dis*. 2006; 65: 415-6.
76. Yılmaz S, Celik G, Esmen SE. Assessment of arterial stiffness in patients with inactive and active behçet's disease. *Scand J Rheumatol*. 2014; 43: 63-9.

P172

ROLE OF ADIPOSE TISSUE AND SKELETAL MUSCLE IN MACROVASCULAR ATHEROSCLEROTIC OCCLUSIVE DISEASE-PERIPHERAL ARTERIAL DISEASE AND CAROTID ARTERY DISEASE

Joana Ferreira
Hospital da Senhora da Oliveira Guimarães, Portugal

Purpose/Background/Objective: Carotid artery disease (CAD) and peripheral arterial disease (PAD) are non-cardiac manifestations of atherosclerotic disease, which are less extensively studied. Presently, adipose tissue (AT) and skeletal muscle (SM) are considered endocrine organs, producing cytokines with vascular effects 1,2,3,4,5,6,7. Main objective is to clarify the role of AT and SM in several stages of atherosclerotic occlusive disease (CAD and PAD), and its connection with arteriosclerotic changes in other territories. Secondary objective: to study the evolution of these parameters after revascularization and after carotid endarterectomy.

Methods: Study Groups (Fig 1): control- without macrovascular atherosclerotic disease eligible for elective varicose veins surgery with inguinal approach. Study group 1- with PAD, confirmed by ankle-brachial index, with or without indication to revascularization. Study group 2- with CAD with or without indication to endarterectomy. We will determine the quantity, endocrine function and histology of SM and AT (in groups submitted to surgery). The groups will be evaluated at admission, 3 and 6 months (Fig 2).

Expected Results: The quantity of AT compartments (visceral, subcutaneous and perivascular) and SM will be determined with transvers abdominal CT scan at the level of 3th vertebra. The endocrine function will be evaluated measuring the myokines and adipokines in blood sample. During the surgery we will collect samples of AT (visceral, subcutaneous and perivascular), SM and artery for histology. We will determine the type, number and size of present cell and vascularization. Additional Central Hemodynamic data will be obtained from carotid Doppler ultrasound, carotid femoral-pulse wave velocity; peripheral central pulse pressure; anthropometric and muscle mass measurements will be performed.

Conclusion: We hope to correlate the atherosclerotic and arteriosclerotic phenotypes with SM and AT characteristics, as well as indexes of sarcopenia.

References

- Jia G, Aroor AR, Soers JR. The role of mineralocorticoid receptor signaling in the cross-talk between adipose tissue and the vascular wall. *Cardiovasc Res*. 2017 Jul 1;113(9):1055-1063.
- Alexandersen P, Tankó LB, Bagger YZ et al. Associations between aortic calcification and components of body composition in elderly Men. *Obesity* 2006;14(9): 1571-8.
- Akoumianakis I, Antoniadis C. The interplay between adipose tissue and the cardiovascular system: is fat always bad? *Cardiovasc Res*. 2017 Jul 1;113(9):999-1008.
- Lau WB, Ohashi K, Wang Y, Ogawa H, Murohara T et al. Role of adipokines in cardiovascular disease. *Circ J*. Jun 23;81(7):920-928.
- Cao ZFH, Cohen ESP. Role of perivascular adipose tissue in vascular physiology and pathology. *Hypertension*. 2017 May;69(5):770-777.
- Guzik TJ, Skiba DS, Touyz RM, Harrison DG. The role of infiltrating immune cells in dysfunctional adipose tissue. *Cardiovas Res*. 2017 Jul 1;113(9):1009-1023.
- Kaysen GA. Association between Inflammation and Malnutrition as Risk Factors of Cardiovascular Disease. *Blood Purif* 2006; 24:51–55.