



## 1.4 Prognostic Relevance of Augmentation Index in Prevalent Cardiovascular Disease and Total Mortality: Data From the General Population

Natalie Arnold<sup>1,\*</sup>, Tommaso Gori<sup>1</sup>, Anika Gündling<sup>1</sup>, Andreas Schulz<sup>1</sup>, Jürgen H. Prochaska<sup>1</sup>, Marina Panova-Noeva<sup>1</sup>, Irene M. Schmidtman<sup>1</sup>, Norbert Pfeiffer<sup>1</sup>, Manfred Beutel<sup>1</sup>, Karl J. Lackner<sup>2</sup>, Thomas Münzel<sup>2</sup>, Philipp S. Wild<sup>1</sup>

<sup>1</sup>University Medical Center of the Johannes Gutenberg-University Mainz, Mainz, Germany

<sup>2</sup>Institute for Clinical Chemistry and Laboratory Medicine of the Johannes Gutenberg-University Mainz, Mainz, Germany

### ABSTRACT

**Objectives:** To investigate the role of peripheral augmentation index (AIx) for the prediction of prevalent cardiovascular disease (CVD) and total mortality in the general population in age- and sex- specific manner.

**Methods:** AIx was measured in 11,250 participants of the population-based Gutenberg Health Study by Endo-PAT device. For analysis, the study population was stratified by age and sex (men: <60/≥60 years; women: <55/≥55 years), since a non-linear increase in AIx with aging with a plateau-building at age of 55 years in women and at 60 years in men was observed.

**Results:** During the 8-yr follow-up, a total of 584 deaths (382 men/202 women) occurred. In males, top tertile (T) of AIx was associated with all-cause mortality (hazard ratio (HR) in younger men was 2.30 (95% CI 1.41–3.76) and 1.42 (95% CI 1.09–1.85) among men ≥60 years (T3 vs T1/T2; adjusted for height, heart rate, CVD risk factors). No association was found in females. A stronger predictive ability was observed in younger males, with AIx values above the reference limit (HR 3.15 (95% CI 1.73–5.74), multivariable-adjusted). Finally, an increased AIx was also associated with almost 2-fold higher prevalence of CVD in males <60 years (Prevalence ratio 1.81 (95% CI 1.33–2.45) for T3 vs T1/T2; or 2.01 (95% CI 1.40–2.86) for AIx values above reference in a fully adjusted model), but not in older males or women.

**Conclusion:** Within the present analysis, AIx demonstrated a strong predictive value for prevalent CVD and worse survival in younger males, thereby underlying its clinical usefulness in this particular population.

© 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).