

Artery Research Vol. **25(S1)**; 2019, p. S66 DOI: https://doi.org/10.2991/artres.k.191224.057; ISSN 1872-9312; eISSN 1876-4401 https://www.atlantis-press.com/journals/artres



P23 Tadalafil Improves Hemodynamic Parameters and Arterial Stiffness in Patients with Grade I-II Obesity Without Comorbidities

Mariana Larios-Cardenas*, Fernando Grover-Páez, David Cardona-Muller, Ernesto Cardona, Oscar I. Gonzalez-Radillo, Jonathan Trujillo-Quiros, Marycruz Barocio-Pantoja, Patricia Quezada-Fernandez, Carlos G. Ramos-Becerra

Physiology Department, University Health Sciences Center, University of Guadalajara, Mexico

ABSTRACT

Background: Obesity is a global health problem, it is associated with diabetes, hypertension, and cardiovascular diseases [1] and negative hemodynamic effects have been observed [2]. Tadalafil has shown an improvement in endothelial function [3] the aim of the present investigation was to evaluate its effects on hemodynamic parameters in patients with grade I-II obesity.

Objective: To evaluate the acute effect of tadalafil on hemodynamic and arterial stiffness parameters.

Methods: A double blind clinical trial, randomized, was carried out in patients with grade I-II obesity. Seventy patients were allocated to receive placebo or a single dose of 20 mg of tadalafil; hemodynamic (Omron HEM 9000, UNEXEF) and arterial stiffness (Omrom VP1000) parameters were determined before and 24 hours after intervention. The values are expressed in mean \pm SD. Wilcoxon rank test and U de Mann-Whitney were applied. p < 0.05 was considerated as statistically significant.

Results: Both groups were comparable in the baseline. After the intervention a statistically significant changes were shown in the tadalafil group in the diastolic blood pressure (DBP), (.001); second systolic shoulder 2 (SYS2), (.003); augmentation index (Aix), (.049); and braquial-ankle pulse wave velocity (BAPWV), (.005). After that, changes between groups were analyzed and only DBP (-2.39, p = .017) and BAPWV (p = .028) were statistically significant. No changes were observed in flow mediated dilatation (FMD).

 $\textbf{Conclusion:} \ \ \text{Tadalafil modifies DBP and BAPWV but it had not effect on FMD in the acute administration.}$

REFERENCES

- [1] Manuel Moreno G. Definición y clasificación de la obesidad. Revista Médica Clínica Las Condes 2012;23:124-8.
- [2] Contreras-Leal ÉA, Santiago-García J. Obesidad, síndrome metabólico y su impacto en las enfermedades cardiovasculares. Revista biomédica 2011;22:103–15.
- [3] Bryce Moncloa A, Morales-Villegas EC, Urquiaga Calderón J, Larrauri-Vigna C. Recuperando la función endotelial. An Fac Med 2014;75:367–74.
- © 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (http://creativecommons.org/licenses/by-nc/4.0/).