



P68 Usefulness of an Optimal Cut-off in Central Augmentation Pressure for the Detection of Left Ventricular Hypertrophy in Men

Masakazu Obayashi^{1,*}, Shigeki Kobayashi², Hiroataka Yamamoto¹, Yoriomi Hamada¹, Takumi Nanno¹,
Michihiro Kohno³, Masafumi Yano²

¹Sanyo-Onoda City Hospital, Sanyo-Onoda, Japan

²Division of Cardiology, Department of Medicine and Clinical Science, Yamaguchi University Graduate School of Medicine, Ube, Japan

³Kohno Clinic, Hagi, Japan

ABSTRACT

Background: Recently, we showed an age-related increase in augmentation pressure (AP) measured using Mobil-O-Graph (MOG) in normotensive Japanese individuals. However, AP might be a poor index of wave reflection due to the overlap between the forward wave and reflected wave.

Methods: We enrolled untreated hypertensive patients and patients currently on antihypertensive treatment. For 70 patients (median age, 70.5 years; 34 men), M-mode echocardiography was performed for determination of left ventricular hypertrophy (LVH), while hemodynamic measurements were taken using MOG. We investigated the influence of central hemodynamic parameters on LVH.

Results: Spearman correlation coefficients between various parameters [age, height, systolic blood pressure (SBP), mean BP, diastolic BP, central systolic BP (cSBP), and AP] were calculated for LV mass indexed to body surface area (LVMI; g/m²). In men, age ($r = 0.600$, $p = 0.0002$), height ($r = -0.495$, $p = 0.003$), SBP ($r = 0.423$, $p = 0.013$), cSBP ($r = 0.454$, $p = 0.007$), and AP ($r = 0.661$, $p < 0.0001$) were correlated to LVMI. In women, cSBP ($r = 0.334$, $p = 0.044$) and AP ($r = 0.480$, $p = 0.003$) were correlated to LVMI. In men, LVMI ($R^2 = 0.578$, $p = 0.0001$) was significantly associated with AP ($\beta = 1.32 \pm 0.56$, $p = 0.027$) in multivariate regression analysis. In women, no significant independent parameter for LVMI was observed. ROC curve analysis was performed to estimate the utility of AP for the detection of LVH (LVMI >115 g/m²) in men. Area under the ROC curve was 0.83 (95% CI: 0.68–0.99). The optimal cut-off point of 12.5 mmHg produced 79.0% sensitivity and 86.7% specificity.

Conclusion: Higher AP showing >12.5 mmHg calculated by MOG was a significant independent predictor of LVH in male hypertensive patients.

© 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V.
This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).