



P82 Impaired Skin Microvascular Function in Systemic Lupus Erythematosus

Nikolaos Koletsos^{1*}, Eugenia Gkaliagkousi¹, Antonios Lazaridis¹, Panagiota Anyfanti¹, Areti Triantafyllou¹, Konstantina Dipla², Panagiotis Dolgyras¹, Vasiliki Galanopoulou³, Spyros Aslanidis⁴, Stella Douma¹

¹3rd Department of Internal Medicine, Papageorgiou Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece

²Exercise Physiology and Biochemistry Laboratory, Department of Sports Science at Serres, Aristotle University of Thessaloniki, Serres, Greece

³Rheumatology Department, Papageorgiou Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece

⁴Rheumatology Department-2nd Propedeutic Department of Internal Medicine, Hippokraton Hospital, Aristotle University of Thessaloniki, Thessaloniki, Greece

ABSTRACT

Objectives: Alterations in skin microcirculation have been associated with damage in various microvascular beds [1]. Endothelial dysfunction in microcirculation plays a central role in the pathophysiology of rheumatic diseases, contributing substantially to the increased cardiovascular risk of these patients. Laser speckle contrast analysis (LASCA) is a novel non-invasive technique that can be used to evaluate endothelial function of skin microcirculation [2]. Previous studies have shown impaired skin endothelial function in patients with systemic sclerosis. To our knowledge, no previous study has evaluated skin microcirculation in patients with SLE using LASCA [3,4].

Design and Methods: Fifty-two individuals (25 SLE patients and 27 matched controls) were studied. In all subjects, forearm skin blood flow was recorded under standardized conditions using a laser speckle contrast imager (PeriCam PSI NR System, Perimed). Post-occlusive reactive hyperemia (PORH) was assessed following a standardized protocol and data were analyzed with signal processing software (PIMSoft, Perimed). The amplitude of PORH responses was expressed as a percentage increase between peak and baseline perfusion (%).

Results: There were no differences among the two groups in age, sex, body mass index and hypertension status. Post occlusion reperfusion in SLE patients, was significantly lower as compared to non-SLE controls (155.5 ± 53.1 vs $194.5 \pm 40.5\%$ respectively, $p = 0.004$).

Conclusion: Patients with SLE demonstrate impaired skin microvascular endothelial function, providing a link that could explain the increased cardiovascular risk that these patients bear.

REFERENCES

- [1] Nguyen TT, Shaw JE, Robinson C, Kawasaki R, Wang JJ, Kreis AJ, et al. Diabetic retinopathy is related to both endothelium-dependent and -independent responses of skin microvascular flow. *Diabetes Care* 2011;34:1389–93.
- [2] Roustit M, Cracowski JL. Assessment of endothelial and neurovascular function in human skin microcirculation. *Trends Pharmacol Sci* 2013;34:373–84.
- [3] Ruaro B, Sulli A, Alessandri E, Pizzorni C, Ferrari G, Cutolo M. Laser speckle contrast analysis: a new method to evaluate peripheral blood perfusion in systemic sclerosis patients. *Ann Rheum Dis* 2014;73:1181–5.
- [4] Della Rossa A, Cazzato M, d'Ascanio A, Tavoni A, Bencivelli W, Pepe P, et al. Alteration of microcirculation is a hallmark of very early systemic sclerosis patients: a laser speckle contrast analysis. *Clin Exp Rheumatol* 2013;31:S109–S14.

© 2019 Association for Research into Arterial Structure and Physiology. Publishing services by Atlantis Press International B.V. This is an open access article distributed under the CC BY-NC 4.0 license (<http://creativecommons.org/licenses/by-nc/4.0/>).